

Indo-Pacific warming induced by a weakening of the Atlantic Meridional Overturning Circulation

SHANTONG SUN * AND ANDREW F. THOMPSON

Environmental Science and Engineering, California Institute of Technology, Pasadena, California

SHANG-PING XIE

Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California

SHANG-MIN LONG

Key Laboratory of Marine Hazards Forecasting, Ministry of Natural Resources and College of Oceanography, Hohai University, Nanjing, China

ABSTRACT

The reorganization of the Atlantic Meridional Overturning Circulation (AMOC) is often associated with changes in Earth's climate. These AMOC changes are communicated to the Indo-Pacific basins via wave processes and induce an overturning circulation anomaly that opposes the Atlantic changes on decadal to centennial time scales. We examine the role of this transient, inter-basin overturning response, driven by an AMOC weakening, both in an ocean-only model with idealized geometry and in a coupled CO₂ quadrupling experiment, in which the ocean warms on two distinct timescales: a fast decadal surface warming and a slow centennial subsurface warming. We show that the transient inter-basin overturning produces a zonal heat redistribution between the Atlantic and Indo-Pacific basins. Following a weakened AMOC, an anomalous northward heat transport emerges in the Indo-Pacific, which substantially compensates for the Atlantic southward heat transport anomaly. This zonal heat redistribution manifests as a thermal inter-basin seesaw between the high-latitude North Atlantic and the subsurface Indo-Pacific and helps to explain why Antarctic temperature records generally show more gradual changes than the Northern Hemisphere during the last glacial period. In the coupled CO₂ quadrupling experiment, we find that the inter-basin heat transport due to a weakened AMOC contributes substantially to the slow centennial subsurface warming in the Indo-Pacific, accounting for more than half of the heat content increase and sea level rise. Thus, our results suggest that the transient inter-basin overturning circulation is a key component of the global ocean heat budget in a changing climate.

1. Introduction

The reorganization of the Atlantic Meridional Overturning Circulation (AMOC) is typically a key feature of transitions in Earth's climate (Broecker et al. 1985). Changes in the AMOC may impact other aspects of the climate system even outside of the Atlantic basin, with particular attention paid to the coupling between the AMOC and the Southern Ocean in previous studies (*e.g.*, Stocker and Johnsen 2003; Marshall and Speer 2012). Zonal changes in ocean heat content in response to a changing AMOC, on the other hand, have received less attention. Here we propose that changes in the AMOC result in a redistribution of heat between the Atlantic and Indo-Pacific basins through a transient inter-basin overturning circulation, and in the

case of a weakening AMOC, can produce a centennial-scale subsurface warming in the Indo-Pacific (Fig. 1).

Inter-basin pathways that connect the Atlantic and Indo-Pacific Oceans are well established (Gordon 1986; Talley 2013; Newsom and Thompson 2018), but only recently has the transient adjustment of this inter-basin overturning circulation, in response to AMOC changes, been explored (Sun et al. 2020). A weakened AMOC is associated with a reduced formation rate of North Atlantic Deep Water (NADW) that modifies the density structure in the North Atlantic (*e.g.*, Jansen et al. 2018). These changes propagate as Kelvin waves southward along the western boundary, eastward along the equator, and then poleward along the eastern boundary in the Atlantic (Huang et al. 2000; Johnson and Marshall 2004; Cessi et al. 2004), and cause a change in the isopycnal depth on the eastern boundary of the South Atlantic. Thus, a reduction in the AMOC strength is characterized by an anomalous southward,

*Corresponding author address: California Institute of Technology, Pasadena, California
E-mail: shantong@caltech.edu




FIG. 1. Schematic of overturning circulation and heat transport responses to a weakening of the Atlantic Meridional Overturning Circulation (AMOC), assuming no change in atmospheric state. The background color shows potential temperature along 170°W in the Southern Ocean, 20°W in the Atlantic, and 170°W in the Indo-Pacific from the ECCO state estimate (Forget et al. 2015). The thick black dashed lines with arrow represent the overturning circulation anomaly following a weakening of the AMOC. The black hollow arrow shows the heat transport anomaly due to overturning circulation changes. On decadal to centennial timescales, the Indo-Pacific overturning circulation substantially compensates the AMOC changes, leading to a heat transport response with opposing directions in the Atlantic ($H'_{\text{ATL}} < 0$) and Indo-Pacific ($H'_{\text{IP}} > 0$) basins. The inter-basin heat redistribution causes a cooling in the high-latitude North Atlantic and a warming in the subsurface Indo-Pacific. The heat transport into the Southern Ocean, $H'_{\text{SO}} = H'_{\text{ATL}} + H'_{\text{IP}}$, is a small residual of the northern basin transport anomalies on decadal to centennial timescales and slowly intensifies on multi-centennial timescales, based on the MITgcm ocean-only simulations (Section 3).

geostrophically-balanced, upper-ocean volume transport that is established within months. The signal of isopycnal depth changes propagates further eastward around South Africa, via coastal Kelvin waves, into the Indo-Pacific basins, leading to a similar geostrophically-balanced overturning response that is characterized by an anomalous northward, upper-ocean volume transport and a reduced upwelling (*cf.* Sen Gupta et al. 2016). Thus, the Indo-Pacific overturning changes substantially oppose the Atlantic changes (black dashed arrows in Fig. 1).

These overturning circulation perturbations may also impact heat transport and heat content within each basin. The meridional heat transport anomaly (H') caused by overturning circulation changes (ψ') in a basin, defined in Eulerian depth space, can be expressed as

$$H'_{\text{Eulerian}} = -\rho_0 c_p \int_{-z_{\text{bot}}}^0 \frac{\partial \psi'}{\partial z} \bar{\theta} dz. \quad (1)$$

Here ρ_0 is seawater density, c_p is seawater specific heat capacity, $\bar{\theta}$ is the time and zonal-mean potential temperature in the control climate and contributions to the heat transport anomaly due to temperature changes have been neglected (Section 5a), z_{bot} is the depth of the seafloor, and ψ' is the Eulerian-mean overturning circulation streamfunction anomaly. This last term (ψ') is defined as the vertical integral of the meridional velocity anomaly v' , the deviation away from the control climate, also integrated from the western boundary (x_w) to the eastern boundary (x_e). The combination of a weakened AMOC and the vertical distribution of temperature (warmer at the surface) suggests a southward heat transport anomaly in the Atlantic ($H'_{\text{ATL}} < 0$) and a northward heat transport anomaly in the Indo-Pacific ($H'_{\text{IP}} > 0$) (black hollow arrows in Fig. 1). The divergence and convergence of heat transport following a weakening in AMOC strength leads to a cooling in the

high-latitude North Atlantic and a warming in the Indo-Pacific, respectively, suggesting a zonal redistribution of heat between the Atlantic and Indo-Pacific basins by the transient inter-basin overturning circulation (Fig. 1).

Climate models consistently project the AMOC strength to decline in the 21st century due to surface warming and freshening in the North Atlantic (Cheng et al. 2013; Weijer et al. 2020). Many previous AMOC studies have highlighted the impacts local to the Atlantic Ocean (*e.g.*, Xie and Vallis 2012; Winton et al. 2013; Liu et al. 2020). For example, a weakening of the AMOC in a warming climate reduces the northward ocean heat transport in the Atlantic basin, leading to a cooling of the high-latitude North Atlantic that resembles the North Atlantic warming hole (Menary and Wood 2018). This ocean cooling reduces the atmosphere-ocean temperature difference and leads to an enhanced anthropogenic surface heat uptake in the high-latitude North Atlantic ocean (red arrow in Fig. 1) (*e.g.*, Garuba and Klinger 2016; Shi et al. 2018).

In this study, we focus on heat transport and heat content changes that are responses to AMOC perturbations, but occur in regions remote from the Atlantic basin. Specifically, we attribute a slow centennial-scale subsurface warming of the Indo-Pacific to an inter-basin heat transport. Increasing atmospheric CO₂ levels produce ocean warming on two distinct timescales: a fast response, occurring over several years, due to increases in the ocean's mixed layer temperature; and a slow response, occurring over centuries to millennia, due to warming of the deep ocean (Stouffer 2004; Held et al. 2010). By subtracting the global-mean temperature profile after the fast response in a 1% CO₂ experiment, Long et al. (2014) show that the global-mean slow warming is characterized by a prominent subsurface warming. Traditionally, this subsurface warming signature has been attributed to a diffusive mixing that fluxes heat between the surface and the subsurface ocean (*e.g.*, Gregory 2000; Held et al. 2010; Wang et al. 2015), a mechanism that relies on diabatic processes. Instead, we argue that a transient inter-basin overturning circulation, which is largely associated with an adiabatic movement of isopycnals (Sun and Thompson 2020, their Figure 3b), can also contribute substantially to the slow subsurface warming in the Indo-Pacific basins.

Inter-basin exchange may also explain other aspects of heat content changes in individual basins. For example, Shatwell et al. (2020) recently used a suite of idealized climate model simulations to discuss the influence of the AMOC weakening on ocean warming. Their results showed that the Indo-Pacific warms at approximately the same rate as the Atlantic in response to an abrupt CO₂ doubling, despite a larger surface heat uptake rate per unit surface area in the Atlantic basin. Here we suggest that the similar warming rate between the Indo-Pacific and Atlantic is actually due to the heat content redistribution between these two basins by the transient inter-basin over-

turning circulation, driven by the AMOC weakening. This redistribution was not considered in Shatwell et al. (2020).

Modifications to the Atlantic meridional heat transport, associated with transitions in the AMOC, have also been identified as key features of abrupt climate change throughout Earth's history. In particular, these processes have been linked to an oceanic thermal bi-polar seesaw between the North Atlantic and the Southern Ocean that occurs over millennial timescales in glacial and deglacial periods (*e.g.*, Stocker and Johnsen 2003; Barker et al. 2009). Ice core records from Greenland suggest that variability in Northern Hemisphere (NH) climate was dominated by Dansgaard-Oeschger (DO) events throughout the last glacial period. DO events are characterized by large and abrupt fluctuations in temperature in the NH (Dansgaard et al. 1993) that are accompanied by temperature changes of the opposite sign in the Southern Hemisphere (SH). However, in contrast to the abrupt temperature changes in the NH, temperature variations over Antarctica were more gradual and approximately out of phase with the north (Blunier and Brook 2001; Barker et al. 2009). This anti-phasing relationship in temperature changes between Greenland and Antarctica has been linked to changes in the AMOC (*e.g.*, Stocker and Johnsen 2003; Liu et al. 2009). In this paradigm, a surface forcing perturbation in the North Atlantic weakens the AMOC and reduces Atlantic northward heat transport [Eq. (1)], leading to a cooling in the high-latitude North Atlantic and warming in the SH. Stocker and Johnsen (2003) suggested that the Southern Ocean acts as a heat reservoir and the slower change in SH temperature occurs due to a diffusive coupling of the Southern Ocean heat storage with the reduced AMOC. They argued that this gives rise to a 1000 yr time scale for SH changes although this value is empirical in their study. Again, invoking inter-basin heat exchange provides a more physical explanation for this behavior. More concretely, the gradual rate of Antarctica temperature change, in comparison to abrupt NH changes, occurs because much of the reduced northward Atlantic heat transport related to AMOC modifications is compensated by a northward heat transport anomaly in the Indo-Pacific, caused by a transient inter-basin overturning circulation on centennial time scales. The heat transport into the Southern Ocean, which governs the temperature variations recorded in Antarctic ice cores (Markle et al. 2017), arises from the residual between the Atlantic and Indo-Pacific heat transport. This residual is considerably weaker than the basin heat transports and only becomes significant on multi-centennial to millennial timescales, determined by the adjustment timescales of the Southern Ocean overturning circulation (Thompson et al. 2019; Sun et al. 2020).

The goal of this study is to illustrate the magnitude and structure of the heat redistribution carried out by the transient inter-basin overturning circulation and to examine its role in ocean temperature variations in a changing climate.

The structure of the paper is as follows. In Section 2, we introduce two separate model frameworks, ocean-only and coupled simulations, as well as the diagnostic techniques that we apply in this study. We diagnose and explain the inter-basin heat transport due to AMOC changes in Section 3. We compare the dynamics in the ocean-only model to the behavior of the inter-basin exchange in a coupled model and connect the inter-basin heat transport to a slow warming in the Indo-Pacific in response to an increased CO₂ forcing in Section 4. A discussion of the results is provided in Section 5, followed by a brief summary in Section 6.

2. Approach and Methodology

Our approach is to use a hierarchy of numerical models to illustrate the dynamics of inter-basin heat transport and its climate impacts. We start with ocean-only simulations to isolate the relationships between basin-scale overturning circulations and heat transports. Yet, the ocean-only model omits a number of key components of the climate system, including feedbacks between the atmosphere and ocean. To account for these limitations, we next analyze output from a coupled model and assess the relative importance of inter-basin heat transport as compared to surface heat uptake on ocean warming under increased CO₂ radiative forcing.

a. Ocean-only simulations

We use the Massachusetts Institute of Technology General Circulation Model (MITgcm; Marshall et al. 1997) to carry out ocean-only simulations that use a nonlinear equation of state (Jackett and McDougall 1995). The model domain, discretized on a 1°-resolution grid, includes three idealized basins that represent the Atlantic, Indian, and Pacific Ocean, connected in the south with a re-entrant channel that represents the Southern Ocean (Fig. 2a), which is a similar domain to the reduced gravity model in Sun and Thompson (2020). In the vertical, there are 30 vertical layers ranging from 20 m at the top to 250 m at the bottom. A vertical diffusivity is implemented that is a function of depth and increases from $2.0 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ at the surface to $1.0 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$ with a transition depth of 2000 m (Bryan and Lewis 1979). Vertical convection is represented by an implicit diffusion with a diffusivity of $10 \text{ m}^2 \text{ s}^{-1}$, whenever the stratification is hydrostatically unstable. Unresolved eddies are represented using the skew-flux form of the Gent-McWilliams (GM) parameterization with an eddy thickness diffusivity $K_{\text{GM}} = 500 \text{ m}^2 \text{ s}^{-1}$. Isopycnal diffusion due to eddies are represented using the Redi scheme with $K_{\text{Redi}} = 500 \text{ m}^2 \text{ s}^{-1}$ (Redi 1982). The momentum is dissipated via Laplacian viscosity, biharmonic viscosity, and vertical viscosity with coefficients $A_h = 1.0 \times 10^4 \text{ m}^2 \text{ s}^{-1}$,

$A_4 = 1.0 \times 10^{12} \text{ m}^4 \text{ s}^{-1}$, and $A_v = 1.0 \times 10^{-3} \text{ m}^2 \text{ s}^{-1}$, respectively¹. The biharmonic viscosity is used to suppress grid-scale noise.

The model is forced with a zonally-uniform zonal wind stress (blue curve in Fig. 2b). Temperature (θ) at the surface is restored to a prescribed zonally-uniform temperature profile (Fig. 2c) with a relaxation timescale of 20 days (*cf.* Haney 1971). For the salinity surface boundary condition, we use a prescribed freshwater flux F_w (Fig. 2d) that is uniform in the zonal direction except in the Pacific Ocean, where an extra constant freshwater flux of $0.3 \times 10^{-8} \text{ m s}^{-1}$ is added to suppress convection in the high-latitude North Pacific. To avoid possible bi-stability due to the mixed surface boundary conditions (Stommel 1961), we also relax the surface salinity to a constant, 35 g kg^{-1} , on a timescale of 2 years (*cf.* Griffies et al. 2009).

We run the model for 3,000 years, at which point the model reaches an approximate steady state, as determined by changes in the AMOC strength. We take the steady state simulation as a control (“Control”) and then apply a 0.05 Sv freshwater hosing in the North Atlantic (hatched area in Fig. 2a) to perturb the AMOC. The simulation is run for another 250 years (“Hosing”), during which the freshwater hosing is kept constant. We compare the differences in the volume and heat transports as well as heat content between the control and perturbation runs due to freshwater hosing. We note that the hosing experiment is only intended to show the oceanic response to AMOC perturbations, such as in a warming climate or during DO events; the simulations are not designed to reproduce comprehensive climate responses given that the model omits key components of the climate system.

b. Coupled simulations

The AMOC is projected to decline in response to anthropogenic greenhouse gas forcing. In order to evaluate the inter-basin heat redistribution and its impact on basin-scale temperature changes related to a modified AMOC in a future, warming climate, we examine an abrupt CO₂ quadrupling (4xCO₂) experiment. Specifically, we make use of the NCAR Community Climate System Model, version 4 (CCSM4; Gent et al. 2011), which is part of the Coupled Model Intercomparison Project, Phase 5 (CMIP5; Taylor et al. 2012). The CCSM4 4xCO₂ experiment, with realistic geography, is initialized from an approximately equilibrated CCSM4 preindustrial run at year 1850, but with the atmospheric CO₂ level instantaneously quadrupled. Both the preindustrial and the 4xCO₂ simulations are continued from 1850 for another 250 years.

¹Note that the description of the model in Sun et al. (2020) stated $A_h = 1.0 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$, but the actual value used was $A_h = 1.0 \times 10^4 \text{ m}^2 \text{ s}^{-1}$.




FIG. 2. Model configurations for the MITgcm ocean-only simulations. (a) Geometry of the model domain. The ocean is 4000 m deep everywhere except for a 2500 m deep submarine sill in the Southern Ocean that represents Drake Passage. The black shading represents continents. The hatched area indicates the region where the 0.05 Sv freshwater hosing is applied. Surface forcing applied to the model: (b) zonal wind stress τ_x (blue; N m^{-2}), (c) restoring surface temperature θ_s ($^{\circ}\text{C}$), and (d) freshwater flux F_w (m s^{-1}). The orange dashed line in (b) represents a wind stress forcing with southward-shifted Southern Hemisphere westerlies that mimic the CCSM4 abrupt $4\times\text{CO}_2$ experiment, as discussed in Text S2 of the supporting information.




FIG. 3. Isopycnal overturning circulation in the MITgcm simulations, mapped to depth coordinate (see details in Appendix A). (a,b,c) Meridional overturning circulation streamfunction in the Control simulation, averaged over the last 50 years. (d,e,f) Changes in the overturning circulation due to freshwater hosing, calculated as the difference in overturning circulation streamfunction of the Hosing run from Control, averaged over the last 50 years. “S. Ocean” stands for “Southern Ocean.” Positive and negative values indicate clockwise and anticlockwise overturning circulations, respectively.

In the following analysis, we focus on the differences between the preindustrial and $4\times\text{CO}_2$ runs.

c. Model diagnostics

The isopycnal overturning circulation streamfunction Ψ is used to represent the meridional overturning circulation

and to define the overturning-driven heat transport (Appendix A). The streamfunction Ψ , defined in isopycnal coordinates, accounts for contributions due to both transient and standing eddies, and it is considered to be a more accurate description of the overturning circulation than the overturning circulation streamfunction defined in depth space (Viebahn and Eden 2012; Ballarotta et al. 2013), which is typically used in the climate literature. The Eulerian-mean streamfunction [Eq. (A3)], defined in depth space, is generally an acceptable approximation of the overturning circulation in the low and mid-latitudes, but it can significantly misrepresent the overturning circulation in the high-latitudes, including the Southern Ocean and the subpolar North Atlantic, even after accounting for transient eddies. As such, the magnitude of the heat transport associated with the Eulerian-mean overturning circulation [Eq. (1)] is usually smaller than the heat transport associated with the isopycnal overturning circulation [Eq. (A4)].

Changes in the basin-scale heat content (\mathcal{H} ; J) can be attributed to anomalies in three different physical processes (assuming a closed northern boundary): (i) surface heat uptake (H_{sfc}), (ii) lateral heat transport across the southern boundary due to the overturning circulation (H_{OT}), and (iii) additional lateral processes due to gyre circulation and eddy diffusion processes (H_R), *i.e.*,

$$\mathcal{H}'(t) = \int_0^t (H'_{\text{sfc}}(t) + H'_{\text{OT}}(t) + H'_R(t)) dt, \quad (2)$$

where the prime represents a deviation from the control run, and the heat transport terms H have units of W. The estimate of the heat transport by the overturning circulation H_{OT} in Eq. (A4) also includes contributions from the subtropical cell, which is associated with the wind-driven gyre circulation and projects to density/depth space as a shallow overturning circulation (upper 500 m in Fig. 3b, c).

In a changing climate the range of density classes and the ocean's vertical stratification differ substantially between the control and perturbed states, and it becomes challenging to isolate heat transport contributions due to circulation changes in isopycnal coordinate (see further discussion in Section 3 and Section 4). Therefore, calculations in Eulerian depth space are used to determine the evolution of the ocean's vertical temperature distribution due to circulation changes [Eqs. (3) to (7) Section 3b]. We remind the reader that this Eulerian-depth approach typically results in an underestimation of the heat transport associated with the overturning circulation.

3. Inter-basin heat redistribution due to AMOC perturbations

In this section, we present results from the MITgcm ocean-only simulations.

a. Transient inter-basin overturning circulation

In the Control simulation at equilibrium, the overturning circulation in the Atlantic basin is primarily connected to the Southern Ocean through a pole-to-pole overturning circulation (Fig. 3a,b,c). In other words, the majority of NADW, formed in the high-latitude North Atlantic and carried southward by the AMOC, rises to the surface in the Southern Ocean along isopycnals via adiabatic upwelling, where it is transformed to other density classes (Wolfe and Cessi 2011; Marshall and Speer 2012). In contrast, the overturning in the Indo-Pacific basins is mainly characterized by an anti-clockwise cell, associated with a northward flow of Antarctic Bottom Water (AABW) close to the seafloor, transformation into lighter water by interior diapycnal mixing, and a southward return flow at shallower depths.

In response to the 0.05 Sv freshwater hosing perturbation, the AMOC both weakens in strength and shoals in depth (Fig. 3e). The AMOC strength at 30°S, defined as the maximum value of the annual-mean AMOC streamfunction at this latitude, weakens by $\sim 20\%$ over the first 50 years and then remains roughly stable afterward (gray line in Fig. 4a). This 50 year timescale of AMOC weakening is associated with the gyre spin-up/spin-down dynamics in the North Atlantic, in response to the freshwater perturbation, and is determined by the transit time of Rossby waves across the Atlantic basin (Sun et al. 2020). There is also a slow millennial-scale evolution of the overturning circulation that is associated with the adjustment of the deep stratification, which is not resolved in the 250-year simulation (*cf.* Jansen et al. 2018; Sun et al. 2020).

Unlike the steady state balance, the AMOC weakening that occurs during the 250-year simulation is largely decoupled from the Southern Ocean, where the overturning circulation remains relatively unchanged, as found in Sun et al. (2020) (Fig. 3d,e). Instead, AMOC changes are principally compensated by overturning circulation anomalies in the Indo-Pacific basins (Fig. 3e,f). The Indo-Pacific overturning response is characterized by a northward transport anomaly in the upper 1.8 km and a southward transport anomaly in the abyss, leading to a volume flux convergence anomaly in the upper Indo-Pacific that is largely balanced by an adiabatic deepening of the isopycnals (Sun and Thompson 2020). This isopycnal deepening results in a subsurface warming through the heave effect (Purkey and Johnson 2013), as discussed in the next subsection.

We quantify the meridional volume transport across 30°S within the upper 1 km, which carries the upper, northward-flowing branch of the AMOC (gray and blue lines in Fig. 4a), in the Atlantic and the Indo-Pacific basins. The anomalous northward volume transport in the upper Indo-Pacific largely cancels the anomalous southward transport in the Atlantic Ocean. The approximately




FIG. 4. Anomalies in volume and heat transport following a weakening of AMOC strength in the MITgcm ocean-only simulations. (a) Meridional volume transport anomaly (Sv) in the upper 1 km across 30°S in the Atlantic (blue), Indo-Pacific (green), and the sum of these two basins (orange). The AMOC strength anomaly at 30°S , defined as the maximum value of the overturning circulation streamfunction, is plotted as the gray line. (b) Overturning-driven meridional heat transport anomaly (PW) [Eq. (A4)] across 30°S in the Atlantic (blue), Indo-Pacific (green), and the sum of these two basins (orange). The meridional heat transport anomaly due to all processes in the Indo-Pacific is plotted as a green dashed line for comparison. Unless otherwise noted, time series throughout this study are smoothed using a 5-year running-mean to suppress noise.

synchronous, but opposite, responses between the Atlantic and Indo-Pacific are due to the fast propagation of coastal Kelvin waves. The residual between the Atlantic and Indo-Pacific volume transport (orange line in Fig. 4a) denotes changes in the Southern Ocean overturning circulation, which responds on multi-centennial timescales, as discussed in Sun et al. (2020).

b. Inter-basin heat redistribution and subsurface Indo-Pacific warming

The transient, inter-basin overturning circulation response described above is also associated with an inter-basin heat transport that effectively redistributes heat from the high-latitude North Atlantic into the Indo-Pacific basins. This transient redistribution occurs via the same mechanism, but with the opposite sense, that transports heat or buoyancy from the low-latitude Pacific to sites of heat loss and NADW formation in the mean state, discussed in Newsom and Thompson (2018). Associated with the AMOC weakening, there is an anomalous overturning-driven southward heat transport out of the Atlantic basin that peaks at ~ 0.15 PW after about 50 years and then remains roughly constant over the next 200 years (blue line in Fig. 4b). This southward heat transport anomaly leads to a surface cooling and an anomalous surface heat uptake in the high-latitude North Atlantic due to the surface temperature relaxation condition (not shown). The anomalous southward heat transport in the Atlantic basin is largely compensated by a northward overturning-driven heat transport anomaly into the Indo-Pacific basins

across 30°S (green solid line in Fig. 4b), which dominates the total meridional heat transport anomaly by all processes combined (green dashed line in Fig. 4b). This heat transport compensation is around 100% during the first 100 years and then slowly decreases to around 70% at the end of the 250-year simulation. The sum of the meridional heat transport anomalies across 30°S in the the Atlantic and Indo-Pacific leaves a smaller residual that represents the net heat transport anomaly into the Southern Ocean. This residual transport slowly intensifies over the duration of the simulation; it is about a third of the magnitude of the Atlantic heat transport anomaly after 250 years (orange line in Fig. 4b).

Advection of a temperature anomaly by the unperturbed flow does not play a significant role in the inter-basin heat transport that connects the high-latitude North Atlantic and the Indo-Pacific basins, at least in these simulations (see discussion in 5a). The time scale for an advective change of this nature would be a century or longer, depending on the flow characteristics (*cf.* Fig. 4b). Instead, the heat transport anomaly is enacted through changes to the circulation strength and structure, which include a deepening of the stratification or density surfaces. This deepening is initiated by a reduction in NADW formation, which converges mass in the upper Atlantic basin, and is then propagated into the Indo-Pacific basins in the forms of coastal and equatorial Kelvin waves (Sun et al. 2020; Sun and Thompson 2020).

The heat transport convergence into the Indo-Pacific basins by the transient inter-basin overturning circulation warms the Indo-Pacific (Fig. 1), but it requires a careful assessment of the temperature changes to isolate this con-




FIG. 5. Subsurface warming and heat budget of the Indo-Pacific basins in the MITgcm ocean-only simulations. (a) Hovmöller diagram of the temperature anomaly ($\bar{\theta}'$), horizontally-averaged over the Indo-Pacific basins, in the perturbed run. (b) Total heat content change (blue) decomposed into anomalous surface heat uptake (orange), heat transport anomalies across the southern boundary by the overturning circulation (red), and the residual (green) [Eq. (2)]. The red dashed line represents the heat content increase due to overturning circulation changes, estimated in Eulerian depth space [integration of Eq. (1)]. (c) Vertical velocity anomaly [w' in Eq. (7)] in balance with the Indo-Pacific Eulerian-mean overturning circulation changes at 30°S , averaged over the last 50 years. (d) Vertical temperature gradient ($\partial\bar{\theta}_c/\partial z$) horizontally-averaged over the Indo-Pacific basins in the Control simulation. (e) Temperature anomaly in the Indo-Pacific due to vertical advection, integrated over 250 years [Eq. (10); blue], compared to the simulated warming at the end of the 250 year simulation ($\bar{\theta}'$; orange). The gray dotted lines in panels (a) and (e) mark the approximate depth of maximum warming in the perturbed run due to freshwater hosing.

tribution to the total basin-scale warming signal. To do this, we examine the evolution of the vertical temperature distribution, averaged over the Indo-Pacific basins, in Eulerian depth space (Fig. 5a). The potential temperature field (θ) in the model interior evolves following

$$\frac{\partial\theta}{\partial t} + \nabla_h \cdot (\vec{V}_h \theta) + \frac{\partial(w\theta)}{\partial z} = \frac{\partial}{\partial z} \left(\kappa \frac{\partial\theta}{\partial z} \right) + \mathcal{R}, \quad (3)$$

where $\vec{V}_h = (u, v)$ is the horizontal velocity vector, the subscript h represents the horizontal direction, κ is diapycnal diffusivity, which we assume only varies as a function of depth. The first term on the right hand side represents

small-scale diapycnal diffusion while the second term \mathcal{R} represents eddy-induced isopycnal diffusion. Integrating Eq. (3) horizontally over the Indo-Pacific basins and dividing it by the Indo-Pacific basins area (A_{IP}), we obtain the evolution of the basin-averaged temperature profile,

$$\bar{\theta} = \frac{1}{A_{\text{IP}}} \iint_{\text{IP}} \theta \, dx \, dy, \quad (4)$$

which is a function of depth and time and is given by

$$\frac{\partial \tilde{\theta}}{\partial t} = \frac{1}{A_{IP}} \left(\int_{x_w}^{x_e} (v\theta)_{30^\circ S} dx - \iint_{IP} \frac{\partial (w\theta)}{\partial z} dx dy \right) \quad (5)$$

$$+ \frac{\partial}{\partial z} \left(\kappa \frac{\partial \tilde{\theta}}{\partial z} \right) + \tilde{\mathcal{R}}.$$

In the perturbed simulation, changes in both the circulation and the temperature distribution can contribute to the evolution of the temperature anomaly away from the Control simulation, $\tilde{\theta}' = \tilde{\theta}_p - \tilde{\theta}_c$:

$$\frac{\partial \tilde{\theta}'}{\partial t} = \frac{1}{A_{IP}} \int_{x_w}^{x_e} (v'\theta_c + v_c\theta')_{30^\circ S} dx \quad (6)$$

$$- \frac{1}{A_{IP}} \iint_{IP} \frac{\partial (w'\theta_c + w_c\theta')}{\partial z} dx dy$$

$$+ \frac{\partial}{\partial z} \left(\kappa \frac{\partial \tilde{\theta}'}{\partial z} \right) + \tilde{\mathcal{R}}',$$

which is linearized by dropping the nonlinear terms, involving $v'\theta'$ and $w'\theta'$. Here, terms with a subscript c are properties evaluated in the equilibrated Control simulation. Our earlier heat budget analysis suggests that the Indo-Pacific warming is driven by an anomalous northward heat transport across the southern boundary, which is dominated by overturning circulation changes (Figs. 4b and 5b). The contributions to heat transport due to advection of temperature anomalies and the nonlinear terms are smaller than the contribution due to overturning circulation changes, as discussed in Section 5a. Therefore, we drop the temperature advection tendency terms that arise from temperature changes ($v_c\theta'$ and $w_c\theta'$) in the equation for $\tilde{\theta}'$. We further approximate the temperature field in the Control simulation, θ_c , using the basin-averaged temperature profile, $\tilde{\theta}_c(z)$, since the temperature gradient in the Control simulation is mainly in the vertical direction. This also motivates us to neglect the isopycnal diffusion term and to replace w' with the basin-averaged vertical velocity anomaly, \tilde{w}' , which must be in balance with the Eulerian-mean overturning circulation streamfunction anomaly at the southern boundary (ψ'_S) (Fig. 5c), *i.e.*, $\tilde{w}' = -\psi'_S/A_{IP}$. With these simplifications, we obtain a vertical advection-diffusion equation for $\tilde{\theta}'$,

$$\frac{\partial \tilde{\theta}'}{\partial t} \approx \frac{1}{A_{IP}} \left(-\frac{\partial \psi'_S}{\partial z} \tilde{\theta}_c + \frac{\partial \psi'_S \tilde{\theta}_c}{\partial z} \right) + \frac{\partial}{\partial z} \left(\kappa \frac{\partial \tilde{\theta}'}{\partial z} \right) \quad (7)$$

$$= -\tilde{w}' \frac{\partial \tilde{\theta}_c}{\partial z} + \frac{\partial}{\partial z} \left(\kappa \frac{\partial \tilde{\theta}'}{\partial z} \right).$$

This advection-diffusion relationship is not intended to reproduce the MITgcm simulations, but is introduced to illustrate the impact of overturning circulation changes on the basin's temperature evolution. The volume integration of Eq. (7) over the Indo-Pacific, multiplied by $\rho_0 c_p$,

is given by,

$$\frac{\partial \mathcal{T}'}{\partial t} = \rho_0 c_p \int_{-z_{bot}}^0 \left(\frac{\partial \psi'_S \tilde{\theta}_c}{\partial z} - \frac{\partial \psi'_S}{\partial z} \tilde{\theta}_c \right) dz + H'_{sfc} \quad (8)$$

$$= \rho_0 c_p \int_{-z_{bot}}^0 -\frac{\partial \psi'_S}{\partial z} \tilde{\theta}_c dz + H'_{sfc},$$

where

$$\int_{-z_{bot}}^0 \frac{\partial \psi'_S \tilde{\theta}_c}{\partial z} dz = (\psi'_S \tilde{\theta}_c) \Big|_{-z_{bot}}^0 = 0,$$

and the boundary conditions

$$\iint_{IP} \kappa \frac{\partial \tilde{\theta}'}{\partial z} dx dy = \frac{H'_{sfc}}{\rho_0 c_p}, \quad @ z = 0, \quad (9)$$

$$\iint_{IP} \kappa \frac{\partial \tilde{\theta}'}{\partial z} dx dy = 0, \quad @ z = -z_{bot},$$

have been used. Thus, heat content changes in the Indo-Pacific basins (\mathcal{T}') have contributions from the heat transport anomaly arising from perturbations in the overturning circulation [red dashed line in Fig. 5b; Eq. (1)] and surface heat uptake H'_{sfc} (orange line in Fig. 5b).

Following a weakened AMOC, the Indo-Pacific develops an overturning circulation anomaly ($\psi'_S > 0$) that opposes the Atlantic changes, associated with an anomalous downwelling ($w' < 0$) and a warming in the Indo-Pacific that peaks at the around 1 km depth ($-w' \frac{\partial \tilde{\theta}_c}{\partial z} > 0$; Fig. 5c,d, e). To compare the subsurface warming driven by ψ'_S with the temperature evolutions in the MITgcm simulations, we integrate the vertical advection term in Eq. (7) in time over 250 years,

$$\tilde{\theta}' \sim - \int_0^{250 \text{ yr}} w' \frac{\partial \tilde{\theta}_c}{\partial z} dt. \quad (10)$$

We find that $\tilde{\theta}'$, derived from (10), qualitatively reproduces the structure of the simulated subsurface warming (Fig. 5e). Critically, the subsurface warming arises from the combination of the structure of the overturning circulation anomaly and its correlation with the vertical temperature profile (Fig. 5c, d). The warming has a weaker contribution from diapycnal mixing (differences between the blue and orange lines in Fig. 5e), which actually diffuses the laterally-induced subsurface warming vertically and leads to an anomalous surface heat loss (orange line Fig. 5b).

Both the MITgcm simulations and the advection-diffusion model demonstrate that a reduced AMOC would drive a heat transport convergence into the Indo-Pacific across the southern boundary, providing an inter-basin pathway for a slow centennial-scale subsurface warming in the Indo-Pacific basins. This subsurface warming is largely due to an adiabatic deepening of the isopycnals,

driven by the transient inter-basin overturning circulation (Text S1 of the Supporting Information). Rather than a downward diffusion of heat following surface warming, here diapycnal mixing results in an upward heat flux as the subsurface temperature anomaly diffuses upward. This upward flux may significantly modify the surface air-sea heat flux (*e.g.*, orange line in Fig. 5b), implying that the transient inter-basin overturning circulation may contribute to the pattern of ocean surface warming (Xie et al. 2010) as well as the strength of regional radiative feedback in a changing climate (Armour et al. 2013). A further discussion of the inter-basin heat redistribution in the context of a proposed bi-polar seesaw mechanism is provided in Section 5b.

4. Indo-Pacific warming due to AMOC slow-down in a warming climate

The MITgcm ocean-only simulations indicate that an inter-basin heat exchange driven by an AMOC weakening can contribute substantially to the warming of the Indo-Pacific basins. Yet, these ocean-only simulations miss key components of the climate system, including some that may also influence ocean basin heat content. Thus, the robustness of the inter-basin heat redistribution is next tested in a more realistic context – a coupled climate simulation under an abrupt CO₂ quadrupling forcing. In the coupled model, the inter-basin heat transport, driven by the AMOC weakening, also contributes substantially to a slow subsurface warming of the Indo-Pacific basins once the ocean surface mixed layer equilibrates with the atmosphere.

a. Changes in circulation and heat transport

In response to the abrupt CO₂ quadrupling, there is an initial weakening of the AMOC and a southward transport anomaly in the upper Indo-Pacific basins of roughly 3 Sv that is short-lived, decaying away within the first decade of the simulation (Fig. 6a; negative values represent southward transport). These initial, sub-decadal scale responses occur due to adjustments to a southward shift in the position of the Southern Hemisphere westerly wind following the CO₂ quadrupling (*e.g.*, Rugenstein et al. 2016; Menzel and Merlis 2019). Specifically, this shift produces an easterly wind anomaly near 30°S (not shown) that is accompanied by an anomalous southward Ekman transport, which manifests as a strengthening of the shallow overturning circulation. This upper-ocean circulation change accounts for the southward heat transport anomaly across 30°S during the first decades in both the Atlantic and the Indo-Pacific basins (Fig. 6b). After the initial decade, as the circulation adjusts to changes in the surface wind stress curl via the propagation of Rossby waves across the basin, the wind-driven gyre circulation becomes slightly weaker than the preindustrial control state. Subsequently, the gyre does not contribute substantially to changes in the heat

transport (see further discussion in Text S2 of the supporting information).

The focus of the remainder of this section is on the centennial-scale overturning circulation changes and its contribution to the heat budget, *i.e.*, the evolution that follows the initial transient decadal response. As shown in Sun and Thompson (2020), the AMOC strength at 30°S weakens by around 8 Sv (40%) during the first 100 years in response to the CO₂ quadrupling and then recovers by ~4 Sv in the following 150 years (gray line in Fig. 6a). Similar to the MITgcm simulations in Section 3, the Indo-Pacific overturning circulation substantially compensates the AMOC changes (blue and green line in Fig. 6a), leaving the Southern Ocean overturning circulation largely unchanged during this 250-year simulations (orange line in Fig. 6a).

The rate of surface heat uptake by the ocean also changes dramatically during the 250-year simulations. During the first 50 years, the Indo-Pacific basin-averaged surface heat flux anomaly decays from around 9 W m⁻² to 0 W m⁻² (positive indicates surface heat uptake) (Fig. 7a), suggesting an approximate thermal balance between the atmosphere and the ocean surface mixed layer in the Indo-Pacific basins. The Indo-Pacific stands in contrast with the Atlantic Ocean and the Southern Ocean, which have large positive surface heat flux anomalies throughout the 250-year simulations, related to the upwelling of deep waters. The Atlantic surface heat flux anomaly, which is largely localized in the subpolar North Atlantic (not shown), changes due to the AMOC evolution, increasing from 6 W m⁻² to 8 W m⁻² as the AMOC weakens and then decreasing to around 4 W m⁻² at the end of the 250-year simulation as the AMOC recovers (*e.g.*, Shi et al. 2018) (Figs. 6a and 7a). The Southern Ocean surface heat flux anomaly, seemingly independent of the AMOC evolution, slowly decreases from 8 W m⁻² to around 3 W m⁻² within 250 years. The different surface heat uptake rates between the Atlantic and Indo-Pacific basins, and the Southern Ocean in response to the abrupt CO₂ forcing leads to thermal steric height gradients between these basins and, thus, necessitates an inter-basin heat exchange to remove this gradient (Newsom et al. 2021).

Similar to the ocean-only simulations, as the AMOC weakens, starting 20 years after the CO₂ perturbation, the Atlantic develops a southward heat transport anomaly and the Indo-Pacific develops a northward heat transport anomaly across the southern boundaries (blue and green lines Fig. 6b). However, unlike the ocean-only simulations in which the residual meridional heat transport anomaly into the Southern Ocean is close to zero due to a large cancellation between Indo-Pacific and Atlantic anomalies, in the coupled simulations the magnitude of the Indo-Pacific heat transport anomaly is roughly two times as large as the Atlantic heat transport anomaly. This leads to a net northward heat transport anomaly out of the Southern Ocean




FIG. 6. Time series of the volume and heat transport anomalies in response to CO₂ quadrupling in CCSM4, similar to Fig. 4. (a) Meridional volume transport anomaly (Sv) in the upper 1 km across 30°S in the Atlantic (blue), Indo-Pacific (green), and the sum of these two basins (orange). The AMOC strength anomaly at 30°S is plotted as a gray line. (b) Overturning-driven meridional heat transport anomaly (PW) [Eq. (A4)] across 30°S in the Atlantic (blue), Indo-Pacific (green), and the sum of these two basins (orange). The meridional heat transport anomaly by all processes in the Indo-Pacific is plotted as a green dashed line for comparison. The gray dotted line marks year 20, when the slow subsurface warming in Indo-Pacific begins.




FIG. 7. Warming of the ocean basins after CO₂ quadrupling in CCSM4. (a) Basin-averaged surface heat flux anomaly (W m^{-2}) in the Atlantic (orange), Indo-Pacific (blue), and Southern Ocean (green). (b) Basin-averaged surface temperature anomaly ($^{\circ}\text{C}$) in the Atlantic (orange), Indo-Pacific (blue), and Southern Ocean (green). (c) Hovmöller diagram of the temperature anomaly (θ'), horizontally-averaged in the Indo-Pacific basins. (d) Scatter plot of the temperature anomaly at 500 m vs the temperature anomaly at the surface, horizontally-averaged in the Indo-Pacific basins. The color indicates years after CO₂ quadrupling. The black dotted lines in panels (a, b) and the “X” in panel (d) indicate year 20, after which the Indo-Pacific temperature evolution is dominated by a subsurface warming.

(orange line in Fig. 6b), which is required to balance the anomalous positive heat uptake in the Southern Ocean throughout the 250-year simulations (Fig. 7a). Since the volume transport anomalies are of similar magnitude between the Indo-Pacific and the Atlantic, the asymmetry in the meridional heat transport anomaly must be due to the different vertical temperature distribution between these two basins. The vertical temperature gradient is smaller in the Atlantic as compared to the Indo-Pacific because NADW is warmer and saltier than its deep water counterpart in the Indo-Pacific (see details in Section 5c). This inter-basin difference in temperature and salinity properties is not reproduced in the MITgcm ocean-only runs. Changes in the gyre circulation due to the southward shift of the SH westerlies do not strongly impact the heat transport asymmetry (Text S2 of the supporting information).

b. Slow subsurface warming in the Indo-Pacific

In response to the abrupt CO₂ forcing, the Indo-Pacific warms on two distinct time scales (Fig. 7b and d). The Indo-Pacific basin-averaged surface temperature increases by $\sim 2^\circ\text{C}$ in 20 years. This first stage is the “fast warming,” which is mainly confined to the surface mixed layer (Fig. 7c,d). The fast warming is followed by a “slow warming” phase of similar magnitude that occurs over the next ~ 200 years. In contrast to the fast warming phase, the slow warming is characterized by a warming signal that is larger in the subsurface than at the surface (Fig. 7d). We present the vertical structure of the slow warming in the Indo-Pacific by calculating the temperature anomaly with respect to the temperature profile at year 20, following Long et al. (2014). The temperature evolution shows a subsurface warming pattern with temperatures increasing by more than 1°C down to 1.5 km depth within 250 years (Fig. 8a). In previous studies this slow subsurface warming has been attributed to a diffusive (downward) heat flux that transfers heat from the surface into the deep ocean (Held et al. 2010; Long et al. 2014). However, applying the model’s upper-ocean (from the mixed layer base to 1 km) mean diapycnal diffusivity of $2 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$, a rough estimate for the timescale τ at which diffusion affects the temperature at a depth $d = 1 \text{ km}$, $\tau \sim d^2/\kappa$, is around 1500 years. This is substantially longer than the timescale associated with the simulated warming of the deep Indo-Pacific ocean (Fig. 7c) and suggests the need to identify a different process to account for the deep warming. We suggest that the deeper warming is instead due to the inter-basin overturning circulation.

Changes in the subsurface temperature during the slow warming phase peak at around 400 m, much shallower than the overturning circulation responses (Fig. 8c, e). This mainly arises from the sharp decrease in the vertical temperature gradient with depth, although there is an additional contribution from diffusion, which is mainly

confined to the upper ocean as discussed below. We estimate the temperature evolution related to the overturning circulation changes by integrating the advection-diffusion equation [Eq. (7)] in time. The integration of the vertical advection term alone [Eq. (10)] approximately reproduces the slow warming at year 250 below 1.5 km (Fig. 8e), suggesting that the deep warming is driven by the inter-basin overturning circulation. However, this approach substantially underestimates the simulated warming at shallower depths (Fig. 8e). Part of this underestimation is due to the neglected vertical diffusion. We estimate the temperature changes due to vertical diffusion alone, which was highlighted in Held et al. (2010) as the driver of the deep ocean warming, by solving a diffusion equation [*cf.* Eq. (7)],

$$\frac{\partial \tilde{\theta}'}{\partial t} = \frac{\partial}{\partial z} \left(\kappa \frac{\partial \tilde{\theta}'}{\partial z} \right), \quad (11)$$

with the boundary conditions as follows,

$$\begin{aligned} \kappa \frac{\partial \tilde{\theta}'}{\partial z} &= F'_{\text{sfc}}, & @ z = 0, \\ \kappa \frac{\partial \tilde{\theta}'}{\partial z} &= 0, & @ z = -z_{\text{bot}}. \end{aligned} \quad (12)$$

Here, $\rho_0 c_p F'_{\text{sfc}} = H'_{\text{sfc}}/A_{\text{IP}}$ is the Indo-Pacific surface heat flux anomaly, given by the blue line in Fig. 7a, and κ is reported by the model and horizontally averaged over the Indo-Pacific basins. Driven by the anomalous surface heat flux due to CO₂ forcing, the temperature changes, from year 20 to year 250, due to vertical diffusion (orange line in Fig. 8e) also peak at the subsurface, with a similar magnitude as the temperature change due to the advection term (blue line in Fig. 8e). However, this diffusive subsurface warming is confined to a shallower depth, mainly in the upper 1 km. Accounting for both vertical advection and diffusion, *i.e.*, solving the advection-diffusion equation (7) with the boundary conditions (12), we recover a larger subsurface warming that comes closer to reproducing the simulated evolution of the temperature profile (green and red lines in Fig. 8e). The additional underestimated warming in the upper 1 km after accounting for both vertical advection and diffusion, we speculate, is largely due to our calculation of the overturning-driven temperature changes in Eulerian space, as discussed in the heat budget analysis below.

Despite a weak surface heat uptake anomaly after year 50 (Fig. 7a), the Indo-Pacific heat content increases steadily and at approximately the same rate throughout the 250 year simulation (blue line in Fig. 8b). A heat budget analysis shows that roughly 75% of the heat content increase and sea level rise during the slow warming, estimated between year 20 and year 150, is due to the heat transport anomaly across the southern boundary by the overturning circulation (red solid line in Fig. 8b), with




FIG. 8. Subsurface warming and heat budget of the Indo-Pacific basin for the slow warming in CCSM4 (*cf.* Fig. 5). (a) Hovmöller diagram of the temperature anomaly ($\delta\theta'$; in reference to year 20) in the CCSM4 4xCO $_2$ experiment, horizontally-averaged over the Indo-Pacific basins. (b) Heat content changes (blue) due to anomalous surface heat uptake (orange), heat transport anomalies across the southern boundary by the overturning circulation (red), and the residual (green). The heat content increase driven by changes in the Eulerian-mean overturning circulation is given by the red dashed line. The equivalent sea level rise, calculated using a constant thermal expansion coefficient $1.5 \times 10^{-4} \text{ }^{\circ}\text{C}^{-1}$, due to heat content increase is labelled on the right ordinate axis. (c) Vertical velocity anomaly in balance with the Indo-Pacific overturning circulation changes at 30 $^{\circ}$ S, averaged over the 250 year simulation. (d) Vertical temperature gradient horizontally-averaged over the Indo-Pacific basin in the preindustrial (PI) run. (e) Temperature changes during the slow warming phase, from year 20 to 250, due to overturning-driven vertical advection [Eq. (10); blue], vertical diffusion [Eq. (11); orange], and the combination of both vertical advection and diffusion [Eq. (7); green]. The temperature anomaly at year 250, associated with the slow warming (panel a), is plotted as a green line for comparison.

the remaining 25% mainly due to surface heat uptake (orange line in Fig. 8b). The heat transport anomaly by the overturning circulation, estimated in the Eulerian space, is smaller than that estimated in isopycnal coordinate (red dashed line vs red solid line in Fig. 8b), likely accounting for the underestimated subsurface warming by the advection-diffusion equation (green and red lines Fig. 8e).

The above analysis highlights inter-basin heat redistribution, driven by a weakening of the AMOC, as a major source for the continued heat content changes and sea level rise in the Indo-Pacific basins following a fast surface mixed layer warming (Fig. 8b). This dominance of

the inter-basin exchange holds even in a coupled climate model where the surface heat uptake is not constrained by imposed surface boundary conditions. This slow warming, peaking at the subsurface, occurs on centennial timescales as volume converges within the upper ocean and is connected to changes in high-latitude processes that modify the ocean overturning circulation (Thompson et al. 2019; Sun and Thompson 2020).




FIG. 9. Heat transport anomaly across 30°S in the CCSM4 abrupt $4x\text{CO}_2$ experiment in comparison to the preindustrial Control simulation. (a) Heat transport anomaly across 30°S in the Atlantic (black) due to changes in the circulation (blue), changes in temperature (orange), and changes in both (green) [Eq. (13)]. (b) Heat transport anomaly across 30°S in the Indo-Pacific (black) due to changes in the circulation (blue), changes in temperature (orange), and changes in both (green). (c) Heat transport anomaly driven by the Eulerian-mean overturning circulation anomaly [Eq. (1)] in the Atlantic (blue) and Indo-Pacific (orange). The green line represents the heat transport calculated using the Atlantic overturning circulation anomaly applied to the Indo-Pacific temperature profile; the sign of this curve is reversed to more easily compare with the orange line [opposite sign of Eq. (14); see Section 5c for details].

5. Discussion

a. Passive vs active warming

The simulations in this study emphasize the role that circulation changes have on ocean heat transport and the

convergence of heat into certain regions of the ocean, a mechanism that has been referred to as “active warming.” However, changes to the ocean’s temperature distribution can also contribute to heat transport anomalies, even in the absence of a change in the ocean circulation, a mechanism referred to as “passive warming.” In the ocean-only simulations, despite a relaxation boundary condition that can change the net surface heat flux, modifications to the vertical temperature profile due to the subsurface warming do not substantially change the vertical temperature gradient from the control state (Fig. 5a,d), and thus, the heat transport anomaly is predominantly due to the overturning circulation changes. However, in the CCSM4 experiments, due to the larger atmospheric perturbation, the ocean’s temperature fields evolve due to radiative forcing and can feedback on the ocean heat transport. Here we compare the contributions to the meridional heat transport across 30°S due to circulation changes and temperature changes in the Atlantic and Indo-Pacific basins in the CCSM4 $4x\text{CO}_2$ experiment.

The meridional ocean heat transport anomaly across 30°S in the $4x\text{CO}_2$ experiment, relative to the preindustrial (PI) run and including contributions from both the gyre circulations as well as the deeper overturning circulation, can be decomposed into three components: changes in circulation, changes in temperature, and changes in both (e.g., Hu et al. 2020). This can be written as:

$$H' = \rho_0 c_p \int_{-z_{\text{bot}}}^0 \int_{x_w}^{x_e} (v'\theta_{\text{PI}} + v_{\text{PI}}\theta' + v'\theta') dx dz, \quad (13)$$

where primes denote anomalies from the PI run. In both the Atlantic and Indo-Pacific basins, the second and third terms are small and of opposite sign. The reason for the leading order cancellation between these two terms is not immediately obvious. However, this cancellation implies that the heat transport anomaly across 30°S is dominated by circulation changes (Fig. 9a, b), such that the dominant contribution to inter-basin heat exchange in the CCSM4 $4x\text{CO}_2$ experiment is linked to overturning circulation changes caused by the AMOC weakening (Figs. 6b and 9).

Although the results of this study highlight the role of active warming in inter-basin heat exchange, passive warming could be important for the Indo-Pacific heat content changes on certain timescales. Recently, Bronselaer and Zanna (2020) suggested that the projected pattern of heat content changes during the 21st century is dominated by passive ocean heat uptake due to increasing atmospheric CO_2 radiative forcing. This passive warming process appears to manifest in the CCSM4 $4x\text{CO}_2$ experiment mainly during the fast warming stage, when the Indo-Pacific heat content changes are mainly due to local surface heat uptake (Fig. 7). However, beyond the 21st century when atmospheric CO_2 reaches equilibrium, surface ocean heat uptake mainly occurs in the high-latitude

North Atlantic and Southern Ocean, and our results suggest that the Indo-Pacific warming will be predominately due to heat redistribution by the inter-basin overturning circulation.

b. Inter-basin seesaw vs bipolar seesaw

This study has, to this point, largely focused on the implications of Indo-Pacific warming due to future climate change. However, rearrangement of the AMOC and the associated changes in meridional heat transport have also been key features of abrupt climate change events throughout Earth's history. Of particular relevance is the thermal bi-polar seesaw, used to describe a change in heat content of the North Atlantic and the Southern Ocean on multi-centennial to millennial timescales associated with Dansgaard-Oeschger (DO) events (Crowley 1992; Stocker and Johnsen 2003). High-latitude forcing, such as Heinrich events, are believed to interrupt NADW formation, after which the AMOC weakens and a southward heat transport anomaly emerges in the Atlantic basin. This southward heat transport anomaly has been postulated as the reason for anti-correlated temperature changes between Greenland and Antarctica during DO events (Crowley 1992): during periods when Greenland warms, Antarctica cools, and vice versa. The more gradual trends in Antarctic temperature as compared to the abrupt Northern Hemisphere changes are explained in Stocker and Johnsen (2003) by assuming that the Southern Ocean acts as a heat reservoir for the southward heat transport (Fig. 10a). More recently, Buizert et al. (2015) used high-accumulation ice cores to show that changes in Northern and Southern Hemispheres temperature were not synchronous, but rather Northern Hemisphere changes lead Southern Hemisphere changes by roughly 200 years. This study provides strong evidence that inter-hemispheric heat redistribution is carried out by the ocean, rather than atmospheric processes and furthermore that this characteristic lag time scale is independent of the background climate state. However, this study did not offer a mechanistic description of the processes that give rise to the 200-yr timescale, although Thompson et al. (2019) suggest that it could be related to the time needed to adjust the stratification and therefore the ventilation of deep waters in the Southern Ocean.

Here we suggest two important updates to the bipolar seesaw theory. First, over a series of studies (Sun et al. 2020; Sun and Thompson 2020), it has been shown that the initial response to a transition in the AMOC takes place principally in the Indo-Pacific basins, not in the Southern Ocean. This is true for changes in both circulation and heat transport. Second, we suggest an inter-basin seesaw, in which the Indo-Pacific, rather than the Southern Ocean, acts as the heat reservoir for the AMOC-driven heat transport changes (Fig. 10b). For instance, when the AMOC weakens during an abrupt Northern Hemisphere

cooling event, the southward heat transport anomaly in the Atlantic was likely substantially compensated by a northward heat transport anomaly into the Indo-Pacific basins, driven by the transient inter-basin overturning circulation (Fig. 1). This inter-basin heat redistribution leads to a cooling in the high-latitude North Atlantic and a subsurface warming in the Indo-Pacific on centennial timescales, *i.e.*, an inter-basin seesaw. The subsurface warming in the Pacific weakens the upper ocean density stratification and may have additionally contributed to enhanced ventilation in the high-latitude North Pacific during Heinrich events (Walczak et al. 2020).

This transfer of heat into the Indo-Pacific is important for two reasons. First, the Indo-Pacific, at 30°S, dominates the zonal extent of the ocean and for this reason, has a stronger impact on the stratification of and transport into the Southern Ocean – essentially, the Indo-Pacific basins provide the northern boundary condition for the depths of the isopycnals that span the ACC (Nikurashin and Vallis 2012; Thompson et al. 2016; Sun et al. 2018). The anomalous subsurface warming of the Indo-Pacific proceeds changes in the Southern Ocean (Fig. 4a) and provides an efficient pathway for the increased, subsurface heat content in the Indo-Pacific to be transported into the Southern Ocean. Second, the adjustment of the Indo-Pacific basins to changes in the surface forcing that perturbs the AMOC happens over a period of multiple decades to a few centuries. The heat transport anomaly into the Southern Ocean, which arises from a residual between the Atlantic and the Indo-Pacific heat transport anomalies, only slowly increases in magnitude as the Southern Ocean overturning circulation responds on multi-centennial timescales (orange lines in Fig. 4). Thus, an inter-basin seesaw between the Atlantic and Indo-Pacific helps to explain the more gradual changes in Antarctic temperature paleo-records as compared to the abrupt changes in the high-latitude North Atlantic during the DO events. The 200-yr lag of SH temperature responses in the inter-hemispheric phase changes is also consistent with the dynamical processes that enable the inter-basin exchange, although further studies are needed to confirm the relative importance of this mechanism in the presence of other forcing changes, such as a perturbation to the strength or position of SH westerlies.

The transient compensation in meridional heat transport between the Atlantic and Indo-Pacific basins in response to an AMOC weakening has been reported in previous studies (*e.g.*, Pedro et al. 2018), and is implicitly included in the conceptual model studies by Thompson et al. (2019), but the mechanisms that give rise to this compensation were not explored. Here we connect the inter-basin heat transport between the Atlantic and Indo-Pacific basins to a transient inter-basin overturning circulation, following AMOC changes.




FIG. 10. Schematics of the (a) bipolar seesaw and (b) inter-basin seesaw mechanisms. The bipolar seesaw schematic is adapted from Stocker and Johnsen (2003, their Figure 2), with the double-arrow representing heat exchange between the Southern Ocean and South Atlantic due to diffusive processes. In panel (b), the orange arrows represent the heat transport anomaly following a weakened AMOC, with the magnitude represented by the width of the arrow. The anomalous heat transport into the Southern Ocean is the residual between the Atlantic (southward) and the Indo-Pacific (northward) heat transport anomalies, and it is weaker than the heat transport anomalies in either of the two northern basins. This revised view of centennial-scale overturning adjustment suggests that the Indo-Pacific is the key heat reservoir associated with Dansgaard-Oeschger (DO) events during the last glacial period, (*cf.* Stocker and Johnsen 2003; Pedro et al. 2018).

c. Southern Ocean's role in the inter-basin heat transport

There remain outstanding questions regarding the Southern Ocean's role in the inter-basin heat transport. For instance, the CCSM4 coupled simulation shows that the magnitude of the heat transport response is larger in the Indo-Pacific (H'_{IP}) as compared to the Atlantic (H'_{ATL}), associated with the transient inter-basin overturning circulation, *i.e.*, $H'_{SO} = H'_{ATL} + H'_{IP} > 0$ (Fig. 6). This behavior differs from the ocean-only simulations in which $H'_{SO} \leq 0$ (Fig. 4). The inter-basin asymmetry in heat transport in the CCSM4 coupled simulation arises from differences in the vertical temperature gradients between the Atlantic and Indo-Pacific basins. This difference in temperature structure is consistent with observations but not simulated in the MITgcm ocean-only runs. To support this, we calculate the hypothetical overturning-driven heat transport anomaly in the Atlantic basin but imposing a temperature profile that is identical to the Indo-Pacific (θ_{IP}), *i.e.*,

$$H'_{ATL} = - \int_{-z_{bot}}^0 \frac{\partial \psi'_{ATL}}{\partial z} \theta_{IP} dz. \quad (14)$$

Indeed, using the same temperature profile in the Indo-Pacific from the PI control run, the heat transport driven by the Eulerian-mean Atlantic overturning circulation (green line in Fig. 9c; reversed in sign for comparison) is approximately the same as the Indo-Pacific (orange line in

Fig. 9c). Crucially, a positive (northward) value for H'_{SO} is consistent with an anomalous surface heat uptake that occurs in the Southern Ocean in CCSM4 after the CO_2 quadrupling (Fig. 7). Yet, it is not obvious whether the heat transport anomaly into the Southern Ocean should be positive or negative in the case of a reduction in AMOC strength but no substantial change in the Southern Ocean surface forcing.

6. Summary

Changes in Earth's climate are often associated with a reorganization of the global ocean overturning circulation. Recently, Sun et al. (2020) showed that there is a transient overturning compensation between the Atlantic and Indo-Pacific basins, following a perturbation in the AMOC. As the AMOC weakens, the Indo-Pacific develops an opposing overturning circulation anomaly, characterized by an anomalous northward surface transport. In this study, we show that this transient inter-basin overturning circulation can effectively redistribute heat between the Atlantic and the Indo-Pacific basins, leading to a slow centennial subsurface warming in the Indo-Pacific. This process makes the dominant contribution to changes in ocean heat content in these basins, even in cases where atmospheric warming leads to increased surface heat uptake. We postulate that there is an inter-basin seesaw associated with the

transient inter-basin overturning circulation that also helps to explain why Antarctic temperature records during the last glacial period generally show more gradual changes than the abrupt transitions recorded in the Northern Hemisphere.

By examining an abrupt CO₂ quadrupling experiment in a coupled climate model, which presents a fast decadal surface warming and a slow centennial subsurface warming, we show that the inter-basin heat redistribution, driven by a weakening of the AMOC, contributes substantially to the slow subsurface warming and sea level rise in the Indo-Pacific in response to CO₂ forcing (e.g., Long et al. 2020). This inter-basin pathway for the Indo-Pacific warming, which hinges on adiabatic processes, is separate from the diapycnal processes emphasized in previous studies that occur over millennial timescales due to the downward diffusion of heat into the deep ocean (e.g., Held et al. 2010). The subsurface warming driven by the transient inter-basin overturning circulation gives rise to an interior heat anomaly in the ocean that can, over time, diffuse upward and potentially contribute to the pattern of ocean surface warming (Xie et al. 2010), influence regional air-sea heat fluxes and atmospheric circulation (Kang et al. 2020; Molina et al. 2021), and modulate regional radiative feedback (Armour et al. 2013).

Acknowledgments. Without implying their endorsement, we thank Emily Newsom, Laure Zanna, Spencer Jones, and Dave Bonan for helpful discussions. We are also grateful for constructive comments from Christopher Wolfe and two anonymous reviewers that helped to improve this manuscript. SS and AFT were supported by NSF award OCE-1756956 and the David and Lucille Packard Foundation. SL acknowledges support from the National Natural Science Foundation of China (42076208 and 41706026). The computations presented here were conducted in the Resnick High Performance Center, a facility supported by Resnick Sustainability Institute at the California Institute of Technology.

Data Availability Statement

The CCSM4 model output was downloaded from the Climate Data Gateway at NCAR (<https://www.earthsystemgrid.org>). The code for reproducing the MITgcm ocean-only simulations is provided at the online open access repository figshare (doi: 10.6084/m9.figshare.16308111).

APPENDIX A

Definition of the overturning circulation streamfunction and heat transport

The meridional overturning circulation streamfunction in a basin is calculated in σ_2 (potential density referenced

to 2000 dbar) coordinate as

$$\Psi(y, \sigma_2) = \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} \int_{x_w}^{x_e} \int_{-z_{\text{bot}}}^0 v_r(x, y, z, t) \times \mathcal{H}(\sigma_2 - \sigma_2'(x, y, z, t)) dz dx dt, \quad (\text{A1})$$

where $v_r = v_{\text{Eulerian}} + v_{\text{eddy}}$ is the total meridional velocity that includes both the Eulerian-mean flow (v_{Eulerian}) and eddy bolus contribution due to parameterized eddies (v_{eddy}), σ_2' is the potential density field calculated by the model, t_1 and t_2 define the averaging period, and $\mathcal{H}(\sigma_2 - \sigma_2')$ is the Heaviside step function of $\sigma_2 - \sigma_2'$ such that $\Psi(y, \sigma_2)$ represents the northward volume transport above the isopycnal denoted by σ_2 .

For presentation purpose, we map the overturning circulation streamfunction Ψ to depth coordinates using the zonal and time averaged depth of each isopycnal (Fig. 3)

$$\phi(y, \bar{\zeta}(y, \sigma_2)) = \Psi(y, \sigma_2), \quad (\text{A2})$$

where $\bar{\zeta}(y, \sigma_2)$ is the zonal and time mean of the isopycnal depth ζ (Nurser and Lee 2004).

In this study, we also use the Eulerian-mean overturning circulation,

$$\psi(y, z) = \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} \int_{x_w}^{x_e} \int_z^0 v_{\text{Eulerian}}(x, y, z', t) dz' dx dt. \quad (\text{A3})$$

As compared to the isopycnal overturning circulation, the Eulerian-mean overturning circulation misses the eddy bolus contribution (v_{bolus}), as well as the contribution due to standing eddies (Viebahn and Eden 2012).

The meridional heat transport associated with the isopycnal overturning circulation is calculated as

$$H_{\text{OT}}(y) = \rho_0 c_p \int_{\sigma_{\text{min}}}^{\sigma_{\text{max}}} \frac{\partial \Psi(y, \sigma_2)}{\partial \sigma_2} \hat{\theta}(y, \sigma_2) d\sigma_2. \quad (\text{A4})$$

Here $\hat{\theta}$ is the thickness-averaged temperature (Young 2012), defined as

$$\hat{\theta}(y, \sigma_2) = \frac{\int_{t_1}^{t_2} \int_{x_w}^{x_e} \theta(x, y, \zeta, t) \gamma(x, y, \sigma_2, t) dx dt}{\int_{t_1}^{t_2} \int_{x_w}^{x_e} \gamma(x, y, \sigma_2, t) dx dt}, \quad (\text{A5})$$

where ζ is the depth of isopycnal σ_2 and $\gamma = \partial \zeta / \partial \sigma_2$ is the isopycnal thickness.

References

- Armour, K. C., C. M. Bitz, and G. H. Roe, 2013: Time-varying climate sensitivity from regional feedbacks. *J. Clim.*, **26** (13), 4518–4534.
- Ballarotta, M., S. Drijfhout, T. Kuhlbrodt, and K. Döös, 2013: The residual circulation of the Southern Ocean: Which spatio-temporal scales are needed? *Ocean Modell.*, **64**, 46–55.
- Barker, S., P. Diz, M. J. Vautravers, J. Pike, G. Knorr, I. R. Hall, and W. S. Broecker, 2009: Interhemispheric Atlantic seesaw response during the last deglaciation. *Nature*, **457** (7233), 1097–1102.

- Blunier, T., and E. J. Brook, 2001: Timing of millennial-scale climate change in Antarctica and Greenland during the last glacial period. *Science*, **291** (5501), 109–112.
- Broecker, W. S., D. M. Peteet, and D. Rind, 1985: Does the ocean-atmosphere system have more than one stable mode of operation? *Nature*, **315** (6014), 21–26.
- Bronslaer, B., and L. Zanna, 2020: Heat and carbon coupling reveals ocean warming due to circulation changes. *Nature*, **584** (7820), 227–233.
- Bryan, K., and L. Lewis, 1979: A water mass model of the world ocean. *J. Geophys. Res.*, **84** (C5), 2503–2517.
- Buizert, C., and Coauthors, 2015: Precise interglacial phasing of abrupt climate change during the last ice age. *Nature*, **520** (7549), 661.
- Cessi, P., K. Bryan, and R. Zhang, 2004: Global seiching of thermocline waters between the Atlantic and the Indian-Pacific Ocean Basins. *Geophys. Res. Lett.*, **31** (4).
- Cheng, W., J. C. Chiang, and D. Zhang, 2013: Atlantic meridional overturning circulation (AMOC) in CMIP5 models: RCP and historical simulations. *J. Clim.*, **26** (18), 7187–7197.
- Crowley, T. J., 1992: North Atlantic deep water cools the Southern Hemisphere. *Paleoceanogr.*, **7** (4), 489–497.
- Dansgaard, W., and Coauthors, 1993: Evidence for general instability of past climate from a 250-kyr ice-core record. *Nature*, **364** (6434), 218.
- Forget, G., J.-M. Campin, P. Heimbach, C. Hill, R. Ponte, and C. Wunsch, 2015: ECCO version 4: An integrated framework for non-linear inverse modeling and global ocean state estimation. *Geoscientific Model Development*, **8**, 3071–3104, doi:10.5194/gmd-8-3071-2015.
- Garuba, O. A., and B. A. Klinger, 2016: Ocean heat uptake and interbasin transport of the passive and redistributive components of surface heating. *J. Clim.*, **29** (20), 7507–7527.
- Gent, P. R., and Coauthors, 2011: The community climate system model version 4. *J. Clim.*, **24** (19), 4973–4991.
- Gordon, A. L., 1986: Inter-ocean exchange of thermocline water. *J. Geophys. Res. Oceans*, **91** (C4), 5037–5046.
- Gregory, J. M., 2000: Vertical heat transports in the ocean and their effect on time-dependent climate change. *Clim. Dyn.*, **16** (7), 501–515.
- Griffies, S. M., and Coauthors, 2009: Coordinated ocean-ice reference experiments (COREs). *Ocean Modell.*, **26** (1), 1–46.
- Haney, R. L., 1971: Surface thermal boundary condition for ocean circulation models. *J. Phys. Oceanogr.*, **1** (4), 241–248.
- Held, I. M., M. Winton, K. Takahashi, T. Delworth, F. Zeng, and G. K. Vallis, 2010: Probing the fast and slow components of global warming by returning abruptly to preindustrial forcing. *J. Clim.*, **23** (9), 2418–2427.
- Hu, S., S.-P. Xie, and W. Liu, 2020: Global pattern formation of net ocean surface heat flux response to greenhouse warming. *J. Clim.*, **33** (17), 7503–7522.
- Huang, R. X., M. A. Cane, N. Naik, and P. Goodman, 2000: Global adjustment of the thermocline in response to deepwater formation. *Geophys. Res. Lett.*, **27** (6), 759–762.
- Jackett, D. R., and T. J. McDougall, 1995: Minimal adjustment of hydrographic profiles to achieve static stability. *J. Atmos. Ocean. Technol.*, **12** (2), 381–389.
- Jansen, M. F., L.-P. Nadeau, and T. M. Merlis, 2018: Transient versus equilibrium response of the ocean's overturning circulation to warming. *J. Clim.*, **31** (13), 5147–5163.
- Johnson, H. L., and D. P. Marshall, 2004: Global teleconnections of meridional overturning circulation anomalies. *J. Phys. Oceanogr.*, **34** (7), 1702–1722.
- Kang, S. M., S.-P. Xie, Y. Shin, H. Kim, Y.-T. Hwang, M. F. Stuecker, B. Xiang, and M. Hawcroft, 2020: Walker circulation response to extratropical radiative forcing. *Science Advances*, **6** (47), eabd3021.
- Liu, W., A. V. Fedorov, S.-P. Xie, and S. Hu, 2020: Climate impacts of a weakened Atlantic Meridional Overturning Circulation in a warming climate. *Science Advances*, **6** (26), eaaz4876.
- Liu, Z., and Coauthors, 2009: Transient simulation of last deglaciation with a new mechanism for Bølling-Allerød warming. *Science*, **325** (5938), 310–314.
- Long, S.-M., S.-P. Xie, Y. Du, Q. Liu, X.-T. Zheng, G. Huang, K.-M. Hu, and J. Ying, 2020: Effects of ocean slow response under low warming targets. *J. Clim.*, **33** (2), 477–496.
- Long, S.-M., S.-P. Xie, X.-T. Zheng, and Q. Liu, 2014: Fast and slow responses to global warming: Sea surface temperature and precipitation patterns. *J. Clim.*, **27** (1), 285–299.
- Markle, B. R., and Coauthors, 2017: Global atmospheric teleconnections during Dansgaard-Oeschger events. *Nat. Geosci.*, **10** (1), 36–40.
- Marshall, J., A. Adcroft, C. Hill, L. Perelman, and C. Heisey, 1997: A finite-volume, incompressible Navier Stokes model for studies of the ocean on parallel computers. *J. Geophys. Res. Oceans*, **102** (C3), 5753–5766.
- Marshall, J., and K. Speer, 2012: Closure of the meridional overturning circulation through Southern Ocean upwelling. *Nat. Geosci.*, **5** (3), 171–180.
- Menary, M. B., and R. A. Wood, 2018: An anatomy of the projected North Atlantic warming hole in CMIP5 models. *Clim. Dyn.*, **50** (7–8), 3063–3080.
- Menzel, M. E., and T. M. Merlis, 2019: Connecting direct effects of CO₂ radiative forcing to ocean heat uptake and circulation. *J. Adv. Model. Earth Syst.*, **11** (7), 2163–2176.
- Molina, M. J., A. Hu, and G. A. Meehl, 2021: Response of Global SSTs and ENSO to the Atlantic and Pacific Meridional Overturning Circulations. *J. Clim.*, 1–68.
- Newsom, E., A. F. Thompson, J. F. Adkins, and E. D. Galbraith, 2021: A hemispheric asymmetry in poleward ocean heat transport across climates: Implications for overturning and polar warming. *Earth Plan. Sci. Lett.*, **568**, 117033.
- Newsom, E. R., and A. F. Thompson, 2018: Reassessing the Role of the Indo-Pacific in the Ocean's Global Overturning Circulation. *Geophys. Res. Lett.*, **45** (22), 12–422.
- Nikurashin, M., and G. Vallis, 2012: A theory of the interhemispheric meridional overturning circulation and associated stratification. *J. Phys. Oceanogr.*, **42** (10), 1652–1667.
- Nurser, A. G., and M.-M. Lee, 2004: Isopycnal averaging at constant height. Part I: The formulation and a case study. *J. Phys. Oceanogr.*, **34** (12), 2721–2739.
- Pedro, J. B., M. Jochum, C. Buizert, F. He, S. Barker, and S. O. Rasmussen, 2018: Beyond the bipolar seesaw: Toward a process understanding of interhemispheric coupling. *Quaternary Sci. Rev.*, **192**, 27–46.
- Purkey, S. G., and G. C. Johnson, 2013: Antarctic Bottom Water Warming and Freshening: Contributions to Sea Level Rise, Ocean Freshwater Budgets, and Global Heat Gain*. *J. Clim.*, **26** (16), 6105–6122.
- Redi, M. H., 1982: Oceanic isopycnal mixing by coordinate rotation. *J. Phys. Oceanogr.*, **12** (10), 1154–1158.
- Rugenstein, M. A., J. M. Gregory, N. Schaller, J. Sedláček, and R. Knutti, 2016: Multiannual ocean-atmosphere adjustments to radiative forcing. *J. Clim.*, **29** (15), 5643–5659.
- Sen Gupta, A., S. McGregor, E. Van Sebille, A. Ganachaud, J. N. Brown, and A. Santoso, 2016: Future changes to the Indonesian Throughflow and Pacific circulation: The differing role of wind and deep circulation changes. *Geophys. Res. Lett.*, **43** (4), 1669–1678.
- Shatwell, P., A. Czaja, and D. Ferreira, 2020: Ocean heat storage rate unaffected by MOC weakening in an idealized climate model. *Geophys. Res. Lett.*, **47** (16), e2020GL089849.

- Shi, J.-R., S.-P. Xie, and L. D. Talley, 2018: Evolving Relative Importance of the Southern Ocean and North Atlantic in Anthropogenic Ocean Heat Uptake. *J. Clim.*, **31** (18), 7459–7479.
- Stocker, T. F., and S. J. Johnsen, 2003: A minimum thermodynamic model for the bipolar seesaw. *Paleoceanogr.*, **18** (4).
- Stommel, H., 1961: Thermohaline convection with two stable regimes of flow. *Tellus*, **13** (2), 224–230.
- Stouffer, R. J., 2004: Time scales of climate response. *J. Clim.*, **17** (1), 209–217.
- Sun, S., I. Eisenman, and A. L. Stewart, 2018: Does Southern Ocean surface forcing shape the global ocean overturning circulation? *Geophys. Res. Lett.*, **45** (5), 2413–2423.
- Sun, S., and A. F. Thompson, 2020: Centennial changes in the Indonesian Throughflow connected to the Atlantic Meridional Overturning Circulation: the ocean's transient conveyor belt. *Geophys. Res. Lett.*, **47**, e2020GL090615.
- Sun, S., A. F. Thompson, and I. Eisenman, 2020: Transient overturning compensation between Atlantic and Indo-Pacific basins. *J. Phys. Oceanogr.*, **50** (8), 2151–2172.
- Talley, L. D., 2013: Closure of the global overturning circulation through the Indian, Pacific, and Southern Oceans: Schematics and transports. *Oceanography*, **26** (1), 80–97.
- Taylor, K. E., R. J. Stouffer, and G. A. Meehl, 2012: An overview of cmip5 and the experiment design. *Bull. Amer. Meteor. Soc.*, **93** (4), 485–498.
- Thompson, A. F., S. K. Hines, and J. F. Adkins, 2019: A Southern Ocean mechanism for the inter-hemispheric coupling and phasing of the bipolar seesaw. *J. Clim.*, **32** (14), 4347–4365.
- Thompson, A. F., A. L. Stewart, and T. Bischoff, 2016: A Multibasin Residual-Mean Model for the Global Overturning Circulation. *J. Phys. Oceanogr.*, **46** (9), 2583–2604.
- Viebahn, J., and C. Eden, 2012: Standing eddies in the meridional overturning circulation. *J. Phys. Oceanogr.*, **42** (9), 1486–1508.
- Walczak, M. H., and Coauthors, 2020: Phasing of millennial-scale climate variability in the Pacific and Atlantic Oceans. *Science*, **370** (6517), 716–720.
- Wang, G., S.-P. Xie, R. X. Huang, and C. Chen, 2015: Robust warming pattern of global subtropical oceans and its mechanism. *J. Clim.*, **28** (21), 8574–8584.
- Weijer, W., W. Cheng, O. Garuba, A. Hu, and B. Nadiga, 2020: CMIP6 models predict significant 21st century decline of the Atlantic Meridional Overturning Circulation. *Geophys. Res. Lett.*, e2019GL086075.
- Winton, M., S. M. Griffies, B. L. Samuels, J. L. Sarmiento, and T. L. Frölicher, 2013: Connecting changing ocean circulation with changing climate. *J. Clim.*, **26** (7), 2268–2278.
- Wolfe, C. L., and P. Cessi, 2011: The adiabatic pole-to-pole overturning circulation. *J. Phys. Oceanogr.*, **41** (9), 1795–1810.
- Xie, P., and G. K. Vallis, 2012: The passive and active nature of ocean heat uptake in idealized climate change experiments. *Clim. Dyn.*, **38** (3–4), 667–684.
- Xie, S.-P., C. Deser, G. A. Vecchi, J. Ma, H. Teng, and A. T. Wittenberg, 2010: Global warming pattern formation: Sea surface temperature and rainfall. *J. Clim.*, **23** (4), 966–986.
- Young, W. R., 2012: An exact thickness-weighted average formulation of the Boussinesq equations. *J. Phys. Oceanogr.*, **42** (5), 692–707.

Supporting Information for “Indo-Pacific warming induced by a weakening of the Atlantic Meridional Overturning Circulation”

SHANTONG SUN * AND ANDREW F. THOMPSON

Environmental Science and Engineering, California Institute of Technology, Pasadena, California

SHANG-PING XIE

Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California

SHANG-MIN LONG

College of Oceanography, Hohai University, Nanjing, China

Introduction

This supporting information contains two sections of text and two figures. In Text S1, the reduced Indo-Pacific upwelling, driven by a weakening AMOC, is partitioned into contributions due to adiabatic and diabatic processes. In Text S2, the influence of Southern Hemisphere wind changes on heat transport is examined with an additional MITgcm simulation.

Text S1: Adiabatic vs Diabatic processes in the subsurface warming

In response to a weakening AMOC, the Indo-Pacific develops an overturning circulation anomaly that opposes the Atlantic changes, characterized by a reduced Indo-Pacific upwelling (or an anomalous downwelling) that accounts for much of the subsurface warming in the ocean-only simulations. Noting that the salinity contribution to the Indo-Pacific buoyancy change is an order of magnitude smaller than the temperature contribution in these simulations, the subsurface warming largely arises from a deepening of isopycnals. This deepening could either be driven by the Indo-Pacific overturning circulation anomaly or a change in diapycnal mixing due to the evolution of the density stratification, as discussed below.

Following Sun and Thompson (2020), we partition the changes in the Indo-Pacific upwelling (w') into a component due to vertical movement of isopycnals (w'_{isop}) and a component due to changes in diapycnal mixing (w'_{diap}):

$$w' = w'_{\text{isop}} + w'_{\text{diap}}. \quad (\text{S1})$$

Integrating Eq. (S1) over an isopycnal surface, σ_2 , within the Indo-Pacific basin, we have

$$-\Psi'_S(\sigma_2, t) = -\frac{\partial \mathcal{V}(\sigma_2, t)}{\partial t} + \iint_{\sigma_2} w'_{\text{diap}}(x, y, z, t) dx dy, \quad (\text{S2})$$

where Ψ'_S represents the Indo-Pacific overturning circulation anomaly at the southern boundary (30°S), defined in Eq. (A1), and \mathcal{V} represents the volume above the isopycnal σ_2 in the Indo-Pacific.

Dividing Eq. (S2) by the horizontal area of the Indo-Pacific basin, A_{IP} , and rearranging the equation, we have the evolution of the mean isopycnal (σ_2) depth,

$$\tilde{h} = \frac{\mathcal{V}}{A_{\text{IP}}}$$

as,

$$\frac{\partial \tilde{h}(\sigma_2, t)}{\partial t} = \frac{\Psi'(\sigma_2, t)}{A_{\text{IP}}} + \frac{1}{A_{\text{IP}}} \iint_{\sigma_2} w'_{\text{diap}}(x, y, z, t) dx dy. \quad (\text{S3})$$

Therefore, changes in the isopycnal depth (labeled as term I below) is driven by a combination of the volume transport anomaly across the southern boundary (term II) and the change in diapycnal mixing (term III),

$$\underbrace{\tilde{h}(\sigma_2, t) - \tilde{h}(\sigma_2, 0)}_{\text{I}} = \underbrace{\frac{1}{A_{\text{IP}}} \int_0^t \Psi'(\sigma_2, t') dt'}_{\text{II}} + \underbrace{\frac{1}{A_{\text{IP}}} \int_0^t \iint_{\sigma_2} w'_{\text{diap}}(x, y, z, t') dx dy dt'}_{\text{III}}. \quad (\text{S4})$$

In the absence of changes in diapycnal mixing (III), the overturning circulation anomaly at the southern boundary

*Corresponding author address: California Institute of Technology, Pasadena, California
E-mail: shantong@caltech.edu




FIG. S1. Changes in the mean isopycnal depth (blue) in the Indo-Pacific and its contributions due to overturning circulation responses (orange) and changes in diapycnal mixing (green), diagnosed using Eq. (S4). The contribution due to changes in diapycnal mixing is calculated as the difference between the blue and orange lines. Four representative isopycnals are selected with their initial depth indicated in parentheses in the title.

is balanced by an adiabatic movement of the isopycnals. On the other hand, in the absence of overturning circulation responses (II), changes in diapycnal mixing (e.g., due to surface forcing perturbations) can also lead to a change in the isopycnal depth and subsurface warming.

We examine the mean isopycnal depth changes and the relative contribution from overturning circulation responses and changes in diapycnal mixing in the MITgcm simulations following Eq. (S4) (Fig. S1). During the first 50 years after the fresh water hosing, changes in the density stratification are small (Fig. 5a) and contributions to the isopycnal deepening related to modifications in diapycnal mixing can be neglected (Fig. S1). In this case, the subsurface warming is almost exclusively due to an adiabatic deepening of isopycnals driven by the overturning circulation anomaly. As the subsurface warming intensifies beyond the first 50 years, diapycnal mixing slowly diffuses the subsurface heat/buoyancy anomaly vertically, which makes a significant contribution to the warming structure at the end of the 250-year simulation. Over much of upper water column, diapycnal mixing weakens the

isopycnal deepening (Fig. S1a,b,c) and represents a cooling effect (the upper 1.9 km in Fig. 5e). In the abyss, by diffusing the subsurface warming downward, changes in diapycnal mixing strengthens the isopycnal deepening (Fig. S1d) and represents a warming effect (below 1.9 km depth in Fig. 5e). Similar behavior occurs in the CCSM4 coupled CO₂ quadrupling experiments (cf. Sun and Thompson 2020, their Figure 3b), except that changes in diapycnal mixing warms the ocean column due to surface heat uptake as CO₂ increases (Fig. 8e).

Text S2: Influence of Southern Hemisphere wind changes on heat transport

Changes in the overturning circulation and inter-basin heat transport are also influenced by a southward shift in the Southern Hemisphere westerly wind in response to CO₂ quadrupling in the CCSM4 simulation (Fig. 6). This change in the surface wind could modify the strength of the gyre circulation and potentially contribute to inter-basin heat exchange (Menzel and Merlis 2019). Here we




FIG. S2. Changes in the volume and heat transport in the MITgcm “Wind+Hosing” experiment. (a) Meridional volume transport anomaly relative to control by the upper 1 km across 30°S in the Atlantic (blue), Indo-Pacific (green), and the combination of these two basins (orange). (b) Meridional heat transport anomaly relative to control by the overturning circulation [Eq. (A4)] across 30°S. (c) Time integration of the meridional heat transport anomaly, difference between the MITgcm “Wind+Hosing” experiment and the “Hosing” experiment, due to changes in the surface wind.

discuss the influence of the surface wind changes on the overturning circulation and inter-basin heat transport.

We carry out an additional MITgcm simulation (“Wind+Hosing”) that is the same as the perturbed run (“Hosing”) in Section 3 but with modifications to the surface wind anomalies that mimic the CCSM4 4xCO₂ experiment (orange line in Fig. 2b). With a change in the surface wind similar to the CCSM4 4xCO₂ experiment in both structure and magnitude, the barotropic subtropi-

cal circulation changes by a similar magnitude as CCSM4 (not shown). The overturning circulation and meridional heat transport responses resemble the CCSM4 4xCO₂ experiments in the first decade (Fig. S2 and Fig. 6) and are characterized by a southward upper ocean volume transport anomaly and southward heat transport anomaly after year 1 in both the Indo-Pacific and the Atlantic, driven by the southward Ekman transport anomaly. Both the upper ocean volume transport and heat transport anomaly reverse signs in the following years as the circulation adjusts to changes in the surface wind forcing until they are dominated by the AMOC changes. This highlights the direct atmospheric effect of increased CO₂ radiative forcing on short timescales (Menzel and Merlis 2019).

Beyond decadal timescales, however, the wind changes do not appear to substantially affect the overturning circulation and heat transport. We integrate in time the differences in the meridional heat transport between the MITgcm “Wind+Hosing” experiment and the “Hosing” experiment (Fig. S2c) to show the impact of wind changes in the heat budget. At the end of the 250 years simulation, the weakened wind-driven gyre circulation, driven by the surface wind changes, leads to an anomalous heat convergence into the Indo-Pacific of roughly 0.35×10^{23} J and an anomalous heat divergence out of the Atlantic by 0.64×10^{23} J. This heat content change is an order of magnitude smaller than the change due to the inter-basin heat exchange driven by the AMOC weakening (cf. Fig. 5b), which has a magnitude around 8×10^{23} J. Based on these MITgcm simulations, we speculate that changes in the gyre circulation related to changes in the Southern Hemisphere wind forcing do not contribute substantially to the inter-basin heat exchange and the slow subsurface warming in the Indo-Pacific in the CCSM4 simulations.

References

- Menzel, M. E., and T. M. Merlis, 2019: Connecting direct effects of CO₂ radiative forcing to ocean heat uptake and circulation. *J. Adv. Model. Earth Syst.*, **11** (7), 2163–2176.
- Sun, S., and A. F. Thompson, 2020: Centennial changes in the Indonesian Throughflow connected to the Atlantic Meridional Overturning Circulation: the ocean’s transient conveyor belt. *Geophys. Res. Lett.*, **47**, e2020GL090615.